arXiv:1203.1453v3 [math.CA] 22 Jun 2012

HERMITE-HADAMARD TYPE INEQUALITIES FOR FUNCTIONS WHOSE DERIVATIVES ARE (α, m)-CONVEX

İMDAT İŞCAN

ABSTRACT. In this paper several inequalities of the right-hand side of Hermite-Hadamard inequality are obtained for the class of functions whose derivatives in absolutely value at certain powers are (α, m)-convex.Some applications to special means of positive real numbers are also given.

1. INTRODUCTION

Let $f : I \subset \mathbb{R} \to \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a.b \in I$ with $a < b$, then

$$f\left(\frac{a+b}{2}\right) \le \frac{1}{b-a}\int_a^b f(x)dx \le \frac{f(a)+f(b)}{2}. \tag{1.1}$$

This doubly inequality is known in the literature as Hermite-Hadamard integral inequality for convex functions.

In [4] Miheşan introduced the class of (α, m)-convex functions as the following:

The function $f : [0, b] \to \mathbb{R}$, $b > 0$, is said to be (α, m)-convex where $(\alpha, m) \in [0, 1]^2$, if we have

$$f(tx + m(1-t)y) \le t^\alpha f(x) + m(1-t^\alpha)f(y)$$

for all $x, y \in [0, b]$ and $t \in [0, 1]$.

It can be easily that for $(\alpha, m) \in \{(0, 0), (\alpha, 0), (1, 0), (1, m), (1, 1), (\alpha, 1)\}$one obtains the following classes of functions: increasing, α-starshaped, starshaped, m-convex, convex, α-convex.

Denote by $K_m^\alpha(b)$ the set of all (α, m)-convex functions on $[0, b]$ for which $f(0) \le 0$. For recent results and generalizations concerning m-convex and (α, m)-convex functions (see [1, 2, 8, 6, 5, 7, 9, 10]).

In [3] Dragomir and Agarwal established the following result connected with the right-hand side of (1.1).

Theorem 1. *Let $f : I \subset \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I°, where $a, b \in I$ with $a < b$.If $|f'|$ is convex on $[a, b]$, then the following inequality holds:*

$$\left|\frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)dx\right| \le \frac{b-a}{8}\left[|f'(a)| + |f'(b)|\right]. \tag{1.2}$$

Date: February 26, 2012.

2000 *Mathematics Subject Classification.* 26A51, 26D15.

Key words and phrases. (α, m)-convex functions,Hermite-Hadamard's inequality, Hölder's integral inequality.

In [10], the following inequality of Hermite-Hadamard type for (α, m)-convex functions holds:

Theorem 2. *Let* $f : [0, \infty) \to \mathbb{R}$ *be an* (α, m)*-convex function with* $(\alpha, m) \in (0, 1]^2$. *If* $0 \leq a < b < \infty$ *and* $f \in L[a, b]$, *then one has the inequality:*

$$\frac{1}{b-a}\int_a^b f(x)dx \leq \min\left\{ \frac{f(a) + \alpha m f\left(\frac{b}{m}\right)}{\alpha+1}, \frac{f(b) + \alpha m f\left(\frac{a}{m}\right)}{\alpha+1} \right\}. \tag{1.3}$$

In [1] the following Hermite-Hadamard type inequalities for $m-$ and (α, m)-convex functions were obtained.

Theorem 3. *Let* I *be an open real interval such that* $[0, \infty) \subset I$. *Let* $f : I \to \mathbb{R}$ *be a differentiable mapping on* I *such that* $f' \in L[a, b]$, *where* $0 \leq a < b < \infty$. *If* $|f'|^q$ *is* m*-convex on* $[a, b]$, *for some fixed* $m \in (0, 1]$ *and* $q \in (1, \infty)$, *then*

$$\begin{aligned} \left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)dx \right| &\leq \frac{b-a}{4}\left(\frac{q-1}{2q-1}\right)^{\frac{q-1}{q}}\left(\mu_1^{\frac{1}{q}} + \mu_2^{\frac{1}{q}}\right) \\ &\leq \frac{b-a}{4}\left(\mu_1^{\frac{1}{q}} + \mu_2^{\frac{1}{q}}\right) \end{aligned} \tag{1.4}$$

where

$$\begin{aligned} \mu_1 &= \min\left\{ \frac{|f'(a)|^q + m\left|f'\left(\frac{a+b}{2m}\right)\right|^q}{2}, \frac{\left|f'(\frac{a+b}{2})\right|^q + m\left|f'\left(\frac{a}{m}\right)\right|^q}{2} \right\}, \\ \mu_2 &= \min\left\{ \frac{|f'(b)|^q + m\left|f'\left(\frac{a+b}{2m}\right)\right|^q}{2}, \frac{\left|f'(\frac{a+b}{2})\right|^q + m\left|f'\left(\frac{b}{m}\right)\right|^q}{2} \right\}. \end{aligned}$$

Theorem 4. *Let* I *be an open real interval such that* $[0, \infty) \subset I$. *Let* $f : I \to \mathbb{R}$ *be a differentiable mapping on* I *such that* $f' \in L[a, b]$, *where* $0 \leq a < b < \infty$. *If* $|f'|^q$ *is* (α, m)*-convex on* $[a, b]$, *for some fixed* $\alpha, m \in (0, 1]$ *and* $q \in [1, \infty)$, *then*

$$\begin{aligned} \left| \frac{f(a)+f(b)}{2} - \frac{1}{b-a}\int_a^b f(x)dx \right| &\leq \frac{b-a}{2}\left(\frac{1}{2}\right)^{1-\frac{1}{q}} \\ &\times \min\left\{ \left(\nu_1 |f'(a)|^q + m\nu_2 \left|f'(\frac{b}{m})\right|^q\right)^{\frac{1}{q}}, \left(\nu_1 |f'(b)|^q + m\nu_2 \left|f'(\frac{a}{m})\right|^q\right)^{\frac{1}{q}} \right\} \end{aligned} \tag{1.5}$$

where

$$\nu_1 = \frac{1}{(\alpha+1)(\alpha+2)}\left[\alpha + \left(\frac{1}{2}\right)^{\alpha}\right]$$

and

$$\nu_2 = \frac{1}{(\alpha+1)(\alpha+2)}\left[\frac{\alpha^2+\alpha+2}{2} - \left(\frac{1}{2}\right)^{\alpha}\right].$$

The main aim of this paper is to establish new inequalities of Hermite-Hadamard type for the class of functions whose derivatives in absolutely value at certain powers are (α, m)-convex.

2. Inequalities for functions whose derivatives are (α, m)-convex

In order to prove our main resuls we need the following lemma:

Lemma 1. *Let $f : I \subset \mathbb{R} \to \mathbb{R}$ be a differentiable mapping on I°, $a, b \in I$ with $a < b$. If $f' \in L[a,b]$ and $\lambda, \mu \in [0,\infty)$, $\lambda + \mu > 0$, then the following equality holds:*

$$\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu} - \frac{1}{b-a}\int_a^b f(x)dx = \frac{b-a}{\lambda+\mu}\int_0^1 \left[(\lambda+\mu)t-\lambda\right] f'(tb+(1-t)a)dt$$

Proof. integration by parts we have

$$\begin{aligned} I &= \int_0^1 \left[(\lambda+\mu)t-\lambda\right] f'(tb+(1-t)a)dt \\ &= \left[(\lambda+\mu)t-\lambda\right] \frac{f(tb+(1-t)a)}{b-a} \Big|_0^1 - \frac{\lambda+\mu}{b-a}\int_0^1 f(tb+(1-t)a)dt \\ &= \frac{\lambda f(a)+\mu f(b)}{b-a} - \frac{\lambda+\mu}{b-a}\int_0^1 f(tb+(1-t)a)dt \end{aligned}$$

Setting $x = tb + (1-t)a$, and $dx = (b-a)\,dt$ gives

$$I = \frac{\lambda f(a)+\mu f(b)}{b-a} - \frac{\lambda+\mu}{(b-a)^2}\int_a^b f(x)dx.$$

Therefore,

$$\left(\frac{b-a}{\lambda+\mu}\right) I = \frac{\lambda f(a)+\mu f(b)}{\lambda+\mu} - \frac{1}{b-a}\int_a^b f(x)dx$$

which completes the proof. □

The next theorem gives a new refinement of the upper Hermite-Hadamard inequality for (α, m)-convex functions.

Theorem 5. *Let $f : I \subset [0,\infty) \to \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a,b]$, where $a, b \in I^\circ$ with $a < b$. If $|f'|^q$ is (α, m)-convex on $[a,b]$, for some fixed $(\alpha, m) \in (0,1]^2$, $\lambda, \mu \in [0,\infty)$ with $\lambda + \mu > 0$, and $q \geq 1$, then the following inequality holds:*

$$\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu} - \frac{1}{b-a}\int_a^b f(x)dx\right| \leq \frac{b-a}{\lambda+\mu}\left(\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}\right)^{\frac{q-1}{q}} \tag{2.1}$$

$$\times \min\left\{\left(\gamma_1 |f'(b)|^q + m\gamma_2 \left|f'(\frac{a}{m})\right|^q\right)^{\frac{1}{q}},\ \left(\gamma_3 |f'(a)|^q + m\gamma_4 \left|f'(\frac{b}{m})\right|^q\right)^{\frac{1}{q}}\right\}$$

where

$$\gamma_1 = \frac{1}{(\alpha+1)(\alpha+2)}\left[\frac{2\lambda^{\alpha+2}}{(\lambda+\mu)^{\alpha+1}} + (\alpha+1)\mu - \lambda\right],\ \gamma_2 = \frac{\lambda^2+\mu^2}{2(\lambda+\mu)} - \gamma_1,$$

and

$$\gamma_3=\frac{1}{(\alpha+1)(\alpha+2)}\left[\frac{2\mu^{\alpha+2}}{(\lambda+\mu)^{\alpha+1}}+(\alpha+1)\lambda-\mu\right],\ \gamma_4=\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}-\gamma_3.$$

Proof. Suppose that $q=1$. From Lemma 1 and using the (α,m)-convexity $|f'|$, we have

$$\begin{aligned}
&\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right| \\
\leq\ & \frac{b-a}{\lambda+\mu}\int_0^1 |(\lambda+\mu)t-\lambda|\,|f'(tb+(1-t)a)|\,dt \\
\leq\ & \frac{b-a}{\lambda+\mu}\int_0^1 |(\lambda+\mu)t-\lambda|\,[t^\alpha\,|f'(b)|+m(1-t^\alpha)]\left|f'(\frac{a}{m})\right|)dt \\
=\ & \frac{b-a}{\lambda+\mu}\int_0^1 |(\lambda+\mu)t-\lambda|\,t^\alpha\,|f'(b)|+m(1-t^\alpha)\,|(\lambda+\mu)t-\lambda|\left|f'(\frac{a}{m})\right|dt
\end{aligned}$$

We have

$$\begin{aligned}
\int_0^1 |(\lambda+\mu)t-\lambda|\,t^\alpha dt &= \int_0^{\frac{\lambda}{\lambda+\mu}} [\lambda-(\lambda+\mu)t]\,t^\alpha dt+\int_{\frac{\lambda}{\lambda+\mu}}^1 [(\lambda+\mu)t-\lambda]\,t^\alpha dt \\
&= \frac{1}{(\alpha+1)(\alpha+2)}\left[\frac{2\lambda^{\alpha+2}}{(\lambda+\mu)^{\alpha+1}}+(\alpha+1)\mu-\lambda\right]=\gamma_1
\end{aligned}$$

and

$$\int_0^1 |(\lambda+\mu)t-\lambda|\,(1-t^\alpha)\,dt=\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}-\gamma_1=\gamma_2,$$

hence

$$\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right|\leq\frac{b-a}{\lambda+\mu}\left(\gamma_1\,|f'(b)|+m\gamma_2\left|f'(\frac{a}{m})\right|\right).$$

Since

$$\int_0^1 |(\lambda+\mu)t-\lambda|\,|f'(tb+(1-t)a)|\,dt=\int_0^1 |(\lambda+\mu)t-\mu|\,|f'(ta+(1-t)b)|\,dt$$

Analogously we obtain

$$\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right|\leq\frac{b-a}{\lambda+\mu}\left(\gamma_3\,|f'(a)|+m\gamma_4\left|f'(\frac{b}{m})\right|\right),$$

where

$$\gamma_3=\frac{1}{(\alpha+1)(\alpha+2)}\left[\frac{2\lambda^{\alpha+2}}{(\lambda+\mu)^{\alpha+1}}+(\alpha+1)\lambda-\mu\right] \text{ and } \gamma_4=\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}-\gamma_3$$

which copmletes the proof for this case.

Suppose now that $q\in(1,\infty)$. From Lemma 1 and using the Hölder's integral inequality, we have

$$\begin{aligned}(2.2)\quad & \int_0^1 |(\lambda+\mu)t-\lambda|\,|f'(tb+(1-t)a)|\,dt\\ \le\; & \frac{b-a}{\lambda+\mu}\left(\int_0^1|(\lambda+\mu)t-\lambda|\,dt\right)^{\frac{q-1}{q}}\left(\int_0^1|(\lambda+\mu)t-\lambda|\,|f'(tb+(1-t)a)|^q\,dt\right)^{\frac{1}{q}}\end{aligned}$$

Since $|f'|^q$ is (α,m)-convex on $[a,b]$, we know that for every $t\in[0,1]$

$$(2.3)\qquad \left|f'(tb+m(1-t)\frac{a}{m})\right|^q\le t^\alpha|f'(b)|^q+m(1-t^\alpha)\left|f'(\frac{a}{m})\right|^q.$$

From the inequalities (**??**), (2.2) and (2.3), we have

$$\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right|\le\frac{b-a}{\lambda+\mu}\left(\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}\right)^{\frac{q-1}{q}}\left(\gamma_1|f'(b)|^q+m\gamma_2\left|f'(\frac{a}{m})\right|^q\right)^{\frac{1}{q}}$$

and analogously

$$\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right|\le\frac{b-a}{\lambda+\mu}\left(\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}\right)^{\frac{q-1}{q}}\left(\gamma_3|f'(a)|^q+m\gamma_4\left|f'(\frac{b}{m})\right|^q\right)^{\frac{1}{q}}$$

which completes the proof. □

Corollary 1. *Suppose that all the assumptions of Theorem5 are satisfied,*

1. *In the inequality (2.1) If we choose* $\lambda=\mu$ *, we obtain the inequality in (1.5).*
2. *In the inequality (2.1) If we choose* $\lambda=\mu$, $m=1$, $q=1$ *and* $\alpha=1$ *we obtain the inequality in (1.2).*
3. *In the inequality (2.1) If we choose* $m=\alpha=1$ *we have*

$$\begin{aligned}(2.4)\quad & \left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right|\\ \le\; & \frac{b-a}{\lambda+\mu}\left(\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}\right)^{\frac{q-1}{q}}\\ & \times\min\left\{\left(\gamma_1|f'(b)|^q+\gamma_2|f'(a)|^q\right)^{\frac{1}{q}},\ \left(\gamma_3|f'(a)|^q+\gamma_4|f'(b)|^q\right)^{\frac{1}{q}}\right\}\end{aligned}$$

where

$$\gamma_1=\frac{1}{6}\left[\frac{2\lambda^3}{(\lambda+\mu)^2}+2\mu-\lambda\right],\ \gamma_2=\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}-\gamma_1,$$

and

$$\gamma_3=\frac{1}{6}\left[\frac{2\mu^3}{(\lambda+\mu)^2}+2\lambda-\mu\right],\gamma_4=\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}-\gamma_3.$$

Theorem 6. *Let $f:I\subset[0,\infty)\rightarrow\mathbb{R}$ be a differentiable mapping on I° such that $f'\in L[a,b]$, where $a,b\in I^\circ$ with $a<b$. If $|f'|^q$ is (α,m)-convex on $[a,b]$, for some fixed $(\alpha,m)\in(0,1]^2$, $\lambda,\mu\in[0,\infty)$ with $\lambda+\mu>0$, and $q>1$, then the following inequality holds:*

$$\begin{aligned}\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right| &\leq \frac{b-a}{(\lambda+\mu)^2}\\ &\times\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left[\lambda^2M_1^{\frac{1}{q}}+\mu^2M_2^{\frac{1}{q}}\right]\end{aligned}\tag{2.5}$$

where

$$\begin{aligned}M_1&=\min\left\{\frac{|f'(a)|^q+\alpha m\left|f'\left(\frac{\lambda b+\mu a}{m(\lambda+\mu)}\right)\right|^q}{\alpha+1},\ \frac{\left|f'(\frac{\lambda b+\mu a}{\lambda+\mu})\right|^q+\alpha m\left|f'\left(\frac{a}{m}\right)\right|^q}{\alpha+1}\right\}\\ M_2&=\min\left\{\frac{|f'(b)|^q+\alpha m\left|f'\left(\frac{\lambda b+\mu a}{m(\lambda+\mu)}\right)\right|^q}{\alpha+1},\ \frac{\left|f'(\frac{\lambda b+\mu a}{\lambda+\mu})\right|^q+\alpha m\left|f'\left(\frac{b}{m}\right)\right|^q}{\alpha+1}\right\}\end{aligned}$$

and $\frac{1}{p}+\frac{1}{q}=1.$

Proof. From Lemma 1 and using the Hölder inequality, we have

$$\begin{aligned}&\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right|\\ \leq\ &\frac{b-a}{\lambda+\mu}\left(\int_0^{\frac{\lambda}{\lambda+\mu}}[\lambda-(\lambda+\mu)t]^p\,dt\right)^{\frac{1}{p}}\left(\int_0^{\frac{\lambda}{\lambda+\mu}}|f'(tb+(1-t)a)|^q\ dt\right)^{\frac{1}{q}}\\ &+\frac{b-a}{\lambda+\mu}\left(\int_{\frac{\lambda}{\lambda+\mu}}^1[(\lambda+\mu)t-\lambda]^p\,dt\right)^{\frac{1}{p}}\left(\int_{\frac{\lambda}{\lambda+\mu}}^1|f'(tb+(1-t)a)|^q\ dt\right)^{\frac{1}{q}}\\ \leq\ &\frac{b-a}{\lambda+\mu}\left[\left(\frac{\lambda^{p+1}}{(p+1)(\lambda+\mu)}\right)^{\frac{1}{p}}\left(\frac{\lambda}{\lambda+\mu}M_1\right)^{\frac{1}{q}}+\left(\frac{\mu^{p+1}}{(p+1)(\lambda+\mu)}\right)^{\frac{1}{p}}\left(\frac{\mu}{\lambda+\mu}M_2\right)^{\frac{1}{q}}\right]\\ =\ &\frac{b-a}{\lambda+\mu}\times\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left[\frac{\lambda^2}{\lambda+\mu}M_1^{\frac{1}{q}}+\frac{\mu^2}{\lambda+\mu}M_2^{\frac{1}{q}}\right]\end{aligned}$$

where we use the fact that

$$\begin{aligned}\int_0^{\frac{\lambda}{\lambda+\mu}} [\lambda-(\lambda+\mu)t]^p\,dt &= \frac{\lambda^{p+1}}{(p+1)(\lambda+\mu)},\\ \int_{\frac{\lambda}{\lambda+\mu}}^{1} [(\lambda+\mu)t-\lambda]^p &= \frac{\mu^{p+1}}{(p+1)(\lambda+\mu)}\end{aligned}$$

and by Theorem2 we get

$$\begin{aligned}&\frac{\lambda+\mu}{\lambda}\int_0^{\frac{\lambda}{\lambda+\mu}} |f'(tb+(1-t)a)|^q\ dt\\ =\ &\frac{1}{\frac{\lambda}{\lambda+\mu}(b-a)}\int_a^{\frac{\lambda b+\mu a}{\lambda+\mu}} |f'(x)|^q\ dx\\ \le\ &\min\left\{\frac{|f'(a)|^q+\alpha m\left|f'\left(\frac{\lambda b+\mu a}{m(\lambda+\mu)}\right)\right|^q}{\alpha+1},\ \frac{\left|f'(\frac{\lambda b+\mu a}{\lambda+\mu})\right|^q+\alpha m\left|f'\left(\frac{a}{m}\right)\right|^q}{\alpha+1}\right\},\\ &\frac{\lambda+\mu}{\mu}\int_{\frac{\lambda}{\lambda+\mu}}^{1} |f'(tb+(1-t)a)|^q\ dt\\ =\ &\frac{1}{\frac{\mu}{\lambda+\mu}(b-a)}\int_{\frac{\lambda b+\mu a}{\lambda+\mu}}^{b} |f'(x)|^q\ dx\\ \le\ &\min\left\{\frac{|f'(b)|^q+\alpha m\left|f'\left(\frac{\lambda b+\mu a}{m(\lambda+\mu)}\right)\right|^q}{\alpha+1},\ \frac{\left|f'(\frac{\lambda b+\mu a}{\lambda+\mu})\right|^q+\alpha m\left|f'\left(\frac{b}{m}\right)\right|^q}{\alpha+1}\right\}.\end{aligned}$$

which completes the proof. □

Corollary 2. *Suppose that all the assumptions of Theorem6 are satisfied, in this case:*

1. In the inequality (2.5) if we choose $\lambda=\mu$ and $\alpha=1$ we obtain the inequality in (1.4).
2. In the inequality (2.5) if we choose $m=\alpha=1$ we have

(2.6)
$$\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right| \le \frac{b-a}{(\lambda+\mu)^2}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left[\lambda^2 M_1^{\frac{1}{q}}+\mu^2 M_2^{\frac{1}{q}}\right]$$

where

$$M_1=\frac{|f'(a)|^q+\left|f'\left(\frac{\lambda b+\mu a}{\lambda+\mu}\right)\right|^q}{2},\ M_2=\frac{|f'(b)|^q+\left|f'\left(\frac{\lambda b+\mu a}{\lambda+\mu}\right)\right|^q}{2}$$

Theorem 7. *Let $f : I \subset [0,\infty) \to \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a,b]$, where $a, b \in I^{\circ}$ with $a < b$. If $|f'|^q$ is (α, m)-convex on $[a,b]$, for some fixed $(\alpha, m) \in (0,1]^2$, $\lambda, \mu \in [0,\infty)$ with $\lambda + \mu > 0$, and $q > 1$, then the following inequality holds:*

$$\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right| \leq \frac{b-a}{\lambda+\mu}\left(\frac{\lambda^{p+1}+\mu^{p+1}}{(p+1)(\lambda+\mu)}\right)^{\frac{1}{p}}\left(\frac{1}{\alpha+1}\right)^{\frac{1}{q}}\min\left\{K_1^{\frac{1}{q}}, K_2^{\frac{1}{q}}\right\}, \tag{2.7}$$

where

$$\begin{aligned} K_1 &= |f'(b)|^q + m\alpha\left|f'(\frac{a}{m})\right|^q \\ K_2 &= |f'(a)|^q + m\alpha\left|f'(\frac{b}{m})\right|^q \end{aligned}$$

and $\frac{1}{p}+\frac{1}{q}=1$.

Proof. From Lemma 1 and using the Hölder's integral inequality, we have

$$\begin{aligned} &\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right| \\ \leq\ & \int_0^1 |(\lambda+\mu)t-\lambda|\,|f'(tb+(1-t)a)|\,dt \\ \leq\ & \frac{b-a}{\lambda+\mu}\left(\int_0^1 |(\lambda+\mu)t-\lambda|^p\,dt\right)^{\frac{1}{p}}\left(\int_0^1 |f'(tb+(1-t)a)|^q\,dt\right)^{\frac{1}{q}} \\ \leq\ & \frac{b-a}{\lambda+\mu}\left(\frac{\lambda^{p+1}+\mu^{p+1}}{(p+1)(\lambda+\mu)}\right)^{\frac{1}{p}}\left(\int_0^1 t^{\alpha}|f'(b)|^q + m(1-t^{\alpha})\left|f'(\frac{a}{m})\right|^q dt\right)^{\frac{1}{q}} \\ =\ & \frac{b-a}{\lambda+\mu}\left(\frac{\lambda^{p+1}+\mu^{p+1}}{(p+1)(\lambda+\mu)}\right)^{\frac{1}{p}}\left(\frac{1}{\alpha+1}\right)^{\frac{1}{q}}\left(|f'(b)|^q + m\alpha\left|f'(\frac{a}{m})\right|^q\right)^{\frac{1}{q}}. \end{aligned}$$

Analogously we obtain

$$\begin{aligned} &\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right| \\ \leq\ & \frac{b-a}{\lambda+\mu}\left(\frac{\lambda^{p+1}+\mu^{p+1}}{(p+1)(\lambda+\mu)}\right)^{\frac{1}{p}}\left(\frac{1}{\alpha+1}\right)^{\frac{1}{q}}\left(|f'(a)|^q + m\alpha\left|f'(\frac{b}{m})\right|^q\right)^{\frac{1}{q}}. \end{aligned}$$

□

Corollary 3. *Suppose that all the assumptions of Theorem7 are satisfied, in this case:*

(1) In the inequality (2.7) if we choose $\lambda = \mu$, then the following inequality holds:

$$\begin{aligned}&\left|\frac{f(a)+f(b)}{2}-\frac{1}{b-a}\int_a^b f(x)dx\right| \\ &\leq \frac{b-a}{2}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{\alpha+1}\right)^{\frac{1}{q}}\min\left\{K_1^{\frac{1}{q}},K_2^{\frac{1}{q}}\right\},\end{aligned}$$

(2) In the inequality (2.7) if we choose $m=\alpha=1$, we have

$$\begin{aligned}&\left|\frac{\lambda f(a)+\mu f(b)}{\lambda+\mu}-\frac{1}{b-a}\int_a^b f(x)dx\right| \\ &\leq \frac{b-a}{\lambda+\mu}\left(\frac{\lambda^{p+1}+\mu^{p+1}}{\lambda+\mu}\right)^{\frac{1}{p}}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1}{q}}\left(\left|f'(a)\right|^q+\left|f'(b)\right|^q\right)^{\frac{1}{q}}\end{aligned} \tag{2.8}$$

3. Some applications for special means

Let us recall the following special means of two nonnegative number a,b with $b\geq a$ and $\alpha\in[0,1]$:

(1) The weighted arithmetic mean

$$A_\alpha(a,b):=\alpha a+(1-\alpha)b,\ a,b\geq 0.$$

(2) The unweighted arithmetic mean

$$A(a,b):=\frac{a+b}{2},\ a,b\geq 0.$$

(3) The weighted harmonic mean

$$H_\alpha(a,b):=\left(\frac{\alpha}{a}+\frac{1-\alpha}{b}\right)^{-1},\quad a,b>0.$$

(4) The unweighted harmonic mean

$$H(a,b):=\frac{2ab}{a+b},\quad a,b>0.$$

(5) The Logarithmic mean

$$L(a,b):=\left\{\begin{array}{cc}\frac{b-a}{\ln b-\ln a} & if\ a\neq b\\ b & if\ a=b\end{array}\right.,\quad a,b>0.$$

(6) The p-Logarithmic mean

$$L_p(a,b):=\left\{\begin{array}{cc}\left(\frac{b^{p+1}-a^{p+1}}{(p+1)(b-a)}\right)^{\frac{1}{p}} & if\ a\neq b\\ b & if\ a=b\end{array}\right.,\quad a,b>0,\ p\in\mathbb{Z}\backslash\{-1,0\}.$$

Proposition 1. *Let $a,b\in\mathbb{R}$ with $0<a<b$ and $n\in\mathbb{Z}$, $|n|\geq 2$. Then, we have the following inequality:*

$$\begin{aligned}\left|A_{\frac{\lambda}{\lambda+\mu}}(a^n,b^n)-L_n^n(a,b)\right| &\leq \frac{b-a}{\lambda+\mu}\left(\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}\right)^{\frac{q-1}{q}} \\ &\times|n|\min\left\{\left(\gamma_1 b^{q(n-1)}+\gamma_2 a^{q(n-1)}\right)^{\frac{1}{q}},\ \left(\gamma_3 a^{q(n-1)}+\gamma_4 b^{q(n-1)}\right)^{\frac{1}{q}}\right\}\end{aligned}$$

where

$$\gamma_1=\frac{1}{6}\left[\frac{2\lambda^3}{(\lambda+\mu)^2}+2\mu-\lambda\right],\ \gamma_2=\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}-\gamma_1,$$
$$\gamma_3=\frac{1}{6}\left[\frac{2\mu^3}{(\lambda+\mu)^2}+2\lambda-\mu\right],\gamma_4=\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}-\gamma_3,$$

$\lambda,\mu\in[0,\infty)$ *with* $\lambda+\mu>0$ *and* $q\geq 1$.

Proof. The assertion follows from inequality (2.4) in Corollary 1, for $f:(0,\infty)\to\mathbb{R},\ f(x)=x^n$ □

Proposition 2. *Let* $a,b\in\mathbb{R}$ *with* $0<a<b$ *and* $n\in\mathbb{Z},\ |n|\geq 2$. *Then, we have the following inequality:*

$$\left|A_{\frac{\lambda}{\lambda+\mu}}(a^n,b^n)-L_n^n(a,b)\right|\leq\frac{b-a}{(\lambda+\mu)^2}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}|n|\left[\lambda^2M_1^{\frac{1}{q}}+\mu^2M_2^{\frac{1}{q}}\right]$$

where

$$M_1=A\left(a^{(n-1)q},A_{\frac{\lambda}{\lambda+\mu}}^{(n-1)q}(b,a)\right)$$
$$M_2=A\left(b^{(n-1)q},A_{\frac{\lambda}{\lambda+\mu}}^{(n-1)q}(b,a)\right),$$

$\lambda,\mu\in[0,\infty)$ *with* $\lambda+\mu>0$, *and* $q>1$ *with* $\frac{1}{p}+\frac{1}{q}=1$.

Proof. The assertion follows from inequality (2.6) in Corollary 2, for $f:(0,\infty)\to\mathbb{R},\ f(x)=x^n$ □

Proposition 3. *Let* $a,b\in\mathbb{R}$ *with* $0<a<b$ *and* $n\in\mathbb{Z},\ |n|\geq 2$. *Then, we have the following inequality:*

$$\begin{aligned}&\left|A_{\frac{\lambda}{\lambda+\mu}}(a^n,b^n)-L_n^n(a,b)\right|\\ \leq\ &\frac{b-a}{\lambda+\mu}\left(\frac{\lambda^{p+1}+\mu^{p+1}}{\lambda+\mu}\right)^{\frac{1}{p}}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}|n|\,A^{\frac{1}{q}}\left(a^{(n-1)q},b^{(n-1)q}\right)\end{aligned}$$

$\lambda,\mu\in[0,\infty)$ *with* $\lambda+\mu>0$, *and* $q>1$ *with* $\frac{1}{p}+\frac{1}{q}=1$.

Proof. The assertion follows from inequality (2.8) in Corollary 3, for $f:(0,\infty)\to\mathbb{R},\ f(x)=\frac{1}{x}$ □

Proposition 4. *Let* $a,b\in\mathbb{R}$ *with* $0<a<b$.*Then we have the following inequality:*

$$\begin{aligned}&\left|H_{\frac{\lambda}{\lambda+\mu}}^{-1}(a,b)-L^{-1}(a,b)\right|\\ \leq\ &\frac{b-a}{\lambda+\mu}\left(\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}\right)^{\frac{q-1}{q}}\\ &\times\min\left\{\left(\gamma_1\frac{1}{b^{2q}}+\gamma_2\frac{1}{a^{2q}}\right)^{\frac{1}{q}},\ \left(\gamma_3\frac{1}{a^{2q}}+\gamma_4\frac{1}{b^{2q}}\right)^{\frac{1}{q}}\right\}\end{aligned}$$

where

$$\gamma_1=\frac{1}{6}\left[\frac{2\lambda^3}{(\lambda+\mu)^2}+2\mu-\lambda\right],\ \gamma_2=\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}-\gamma_1,$$

$$\gamma_3=\frac{1}{6}\left[\frac{2\mu^3}{(\lambda+\mu)^2}+2\lambda-\mu\right],\gamma_4=\frac{\lambda^2+\mu^2}{2(\lambda+\mu)}-\gamma_3,$$

$\lambda,\mu\in[0,\infty)$ *with* $\lambda+\mu>0$ *and* $q\geq 1$.

Proof. The assertion follows from inequality (2.4) in Corollary 1, for $f:(0,\infty)\to\mathbb{R},\ f(x)=x^n$ $\square$

Proposition 5. *Let* $a,b\in\mathbb{R}$ *with* $0<a<b$*.Then we have the following inequality:*

$$\left|H_{\frac{\lambda}{\lambda+\mu}}^{-1}(a,b)-L^{-1}(a,b)\right|\leq\frac{b-a}{(\lambda+\mu)^2}\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left[\lambda^2M_1^{\frac{1}{q}}+\mu^2M_2^{\frac{1}{q}}\right]$$

where

$$M_1=H^{-1}\left(a^{2q},A_{\frac{\lambda}{\lambda+\mu}}^{2q}(b,a)\right),\ M_2=H^{-1}\left(b^{2q},A_{\frac{\lambda}{\lambda+\mu}}^{2q}(b,a)\right)$$

$\lambda,\mu\in[0,\infty)$ *with* $\lambda+\mu>0$*, and* $q>1$ *with* $\frac{1}{p}+\frac{1}{q}=1$.

Proof. The assertion follows from inequality (2.6) in Corollary 2, for $f:(0,\infty)\to\mathbb{R},\ f(x)=\frac{1}{x}$ $\square$

Proposition 6. *Let* $a,b\in\mathbb{R}$ *with* $0<a<b$ *. Then we have the following inequality:*

$$\begin{aligned}&\left|H_{\frac{\lambda}{\lambda+\mu}}^{-1}(a,b)-L^{-1}(a,b)\right|\\ \leq\ &\frac{b-a}{\lambda+\mu}A_{\frac{\lambda}{\lambda+\mu}}^{\frac{1}{p}}(\lambda^p,\mu^p)\left(\frac{1}{p+1}\right)^{\frac{1}{p}}\left(\frac{1}{2}\right)^{\frac{1}{q}}H_{\frac{1}{2}}^{\frac{-1}{q}}\left(a^{2q},b^{2q}\right)\end{aligned}$$

$\lambda,\mu\in[0,\infty)$ *with* $\lambda+\mu>0$*, and* $q>1$ *with* $\frac{1}{p}+\frac{1}{q}=1$.

Proof. The assertion follows from inequality (2.8) in Corollary 3, for $f:(0,\infty)\to\mathbb{R},\ f(x)=\frac{1}{x}$ $\square$

REFERENCES

[1] Bakula, M.K., Ozdemir, M.E. and Pecaric, J. *Hadamard type inequalities for* m*-convex and* (α,m)*-convex functions*, J. Inequal. Pure Appl. Math. 9, Article 96, 2008. [Online: http://jipam.vu.edu.au].

[2] Bakula, M.K., Pecaric, J. and Ribičić, M. *Companion inequalities to Jensen's inequality for* m*-convex and* (α,m)*-convex functions,* J. Inequal. Pure Appl. Math. 7, Article 194, 2006. [Online: http://jipam.vu.edu.au].

[3] Dragomir, S.S. and Agarwal, R.P. *Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula*, Appl. Math. Lett., 11 91-95, 1998.

[4] Miheşan, V.G. *A generalization of the convexity, Seminar on Functional Equations*, Approx. and Convex, Cluj-Napoca, Romania, 1993.

[5] Ozdemir, M.E., Kavurmacı, H. and Set, E. *Ostrowski's type inequalities for* (α,m)*-convex functions*, Kyungpook Math. J. 50, 371-378, 2010.

[6] Ozdemir, M.E., Avcı, M., Set,E. *On some inequalities of Hermite-Hadamard type via* m*-convexity,* Applied Mathematics Letters 23, 1065-1070, 2010.

[7] Ozdemir, M.E., Set, E .and Sarıkaya, M.Z. *Some new Hadamard's type inequalities for co-ordinated* m*-convex and* (α,m)*-convex functions,* Hacettepe Journal of Mathematics and Statistics volume 40 (2), 219-229, 2011.

[8] Ozdemir, M.E., Avcı, M. and Kavurmacı, H. *Hermite-Hadamard-type inequalities via* (α,m)*-convexity,* Computers and Mathematics with Applications 61, 2614-2620, 2011.

[9] Sarıkaya, M.Z., Ozdemir, M.E. and Set, E. *Inequalities of Hermite-Hadamard's type for functions whose derivative's absolute values are m-convex,* RGMIA Res. Rep. Coll. 13, Supplement, Article 5, 2010.

[10] Set,E., Sardari, M.Ozdemir, M.E. and Rooin, J. *On generalizations of the Hadamard inequality for (α, m)-convex functions,* RGMIA Res. Rep. Coll. 12(4), Article 4, 2009.

Department of Mathematics, Faculty of Science and Arts,, Giresun University, 28100, Giresun, Turkey.

E-mail address: `imdat.iscan@giresun.edu.tr`